

# FRONTOTEMPORAL DEMENTIA

Dementia (also known as Neurocognitive Disorder) causes gradual deterioration in attention, decision-making, memory and learning, language, perception and /or social behaviour.

## What is frontotemporal dementia?

Frontotemporal dementia (FTD) is an umbrella term for a group of disorders affecting the frontal and temporal lobes of the brain and is the cause of at least 5% of all dementia cases, possibly more since it is difficult to diagnose. The frontal and temporal regions of the brain can be affected by other forms of dementia. For example, Alzheimers disease can start in the frontal region and poor blood supply to these lobes can occur in vascular dementia. However, generally there are three recognised forms of frontotemporal dementia.

1. **Behavioural variant FTD.** In this condition there are early changes in behaviour, personality and executive function, that is, poor reasoning, judgement and decision-making. The behaviour and personality changes may include apathy, disinhibition (e.g. tactless, socially-inappropriate behaviour), loss of empathy and social interest. The person may develop repeated, pointless movements or sayings. Eating patterns change and the person shows a tendency to always putting something in their mouth (hyperorality).the person is not usually aware of problems and may make disastrous personal or financial decisions due to poor judgement and loss of inhibitions.
2. **Language variants of FTD** There are two sorts of FTD that initially affect language, called *primary progressive aphasia*.
  - (a) In the *semantic* variant the first symptoms are a decline in language skills so that the person has trouble producing words, naming things and structuring sentences. People may struggle to find and understand and pronounce words, especially the ones they do not use very often. The meaning of words can be lost, but grammar remains intact.

- (b) *Non-fluent variation* of primary progressive aphasia: In this condition, speech is slow, laboured and halting and words may be left out and grammar misused. People might have trouble understanding complex sentences. People can retain their writing skills for a long time, but spelling and reading can be impaired.

## Overlap with movement disorders

Perhaps 40% of all people with FTD develop mild motor neurone symptoms such as weakness, stiffness, difficulties with speech and swallowing and changes in emotional expression. About 20% get Parkinson's-like symptoms. Others experience eye movement problems, instability standing and falls.

## Causes

Under the microscope there is loss of brain cells, scarring, and tiny holes in the frontotemporal brain tissues. They may be deposits of abnormal protein especially forms of tau including Pick bodies (named after their discoverer Arnold Pick).

Why these changes occur in the brain is unknown

## Who gets it?

FTD is a common cause of dementia in people younger than 65, only about 25% of cases occur in people over 65 (DSM V). About 40% of people with FTD have a family history of early onset dementia and some specific genes have been identified.

## Diagnosis:

History: The diagnosis is made in the usual manner by the doctor asking the person and other people who know him/her about the onset and course of symptoms. It is important to find out if there is a family history of similar problems. Because it is often difficult to tell FTD from common psychiatric disorders, a GP might ask for a psychiatrist's opinion.

**Laboratory and radiology:** This will test for other causes of impairment. A CT or MRI may show specific changes confirming FTD or another form of dementia

**Neuropsychological testing:** This can indicate abnormalities in the frontal lobe or elsewhere. If there are mainly memory and spatial perception defects and social appropriateness is maintained then the diagnosis is more likely to be Alzheimers disease, affecting the frontotemporal regions.

### How does FTD progress?

FTD is progressive, spreading to the rest of the brain eventually looking like other forms of dementia. The rate of decline tends to be faster than the other dementias with a shorter survival time after diagnosis.

### Treatment:

There is no specific treatment for FTD except for addressing the symptoms.

Treatment with cholinesterase inhibitors is not beneficial in any of these conditions and may worsen behavioural symptoms. Memantine makes no difference to the progression of FTD symptoms.

Certain antidepressants, selective serotonin re-uptake inhibitors (SSRIs) can reduce the severity of compulsions, aggression, abnormal eating, impulsiveness and agitation.

Sometime low doses of atypical antipsychotics help some of the behaviour abnormalities but need to be used cautiously.

---

### References:

American Psychiatric Association (2013) Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM 5). Arlington VA.

Bang, J. Spina, S. & Miller, B. (2015). Frontotemporal dementia. Lancet; 386 1672-82

This publication provides a general summary only of the subject matter covered. People should seek professional advice about their specific case.

**Dementia New Zealand offers support, information and education.  
Ring 0800 4 DEMENTIA or  
0800 433 636.  
Or visit our website at [www.dementia.nz](http://www.dementia.nz)**